

WART THERAPY - CANTHACUR- PS

1. A STRONG BLISTERING AGENT HAS BEEN APPLIED UNDER TAPE OCCLUSION TO YOUR WART(S).
2. PLEASE REMOVE THE APPLIED TAPE AFTER _____ HOURS. REMOVE THE MEDICINE WITH ACETONE OR NAIL POLISH REMOVER.
3. OVER THE NEXT 12-24 HOURS, A BLISTER OR EVEN A BLOOD BLISTER MAY OCCUR. RED INFLAMED SKIN WILL BE UNDERNEATH AND AROUND THE BLISTER, AND THE TREATED AREA WILL BE QUITE TENDER FOR 2-3 DAYS.
4. IF THE BLISTER IS VERY INTENSE AND SORE, YOU MAY CLEANSE THE BLISTER WITH ALCOHOL AND CAREFULLY NICK IT WITH A STERILIZED NEEDLE TO ALLOW BLISTER FLUID TO BE DRAINED. LEAVE THE BLISTER ROOF INTACT.
5. KEEP THE AREA CLEAN WITH PEROXIDE OR ALCOHOL AND THEN APPLY POLYSPORIN OR OTHER ANTIBIOTIC OINTMENT ONE TO TWO TIMES PER DAY AND KEEP THE AREA COVERED WITH A BAND-AID AS NECESSARY.
6. IF YOU HAVE ANY QUESTIONS ABOUT ANY UNUSUAL REACTION AT OR AROUND THE TREATED SITE, PLEASE CONTACT OUR OFFICE.
7. PLEASE KEEP YOUR FOLLOW-UP APPOINTMENT FOR ONGOING THERAPY FOR THE MOST RAPID EFFECTIVE RESPONSE.